

Ab initio study of the vapour-liquid critical point of a symmetrical binary fluid mixture

O V Patsahan, M P Kozlovskii and R S Melnyk §

Institute for Condensed Matter Physics, 1 Svientsitskii Street, Lviv 290011, Ukraine

Abstract. A microscopic approach to the investigation of the behaviour of a symmetrical binary fluid mixture in the vicinity of the vapour-liquid critical point is proposed. It is shown that the problem can be reduced to the calculation of the partition function of a 3D Ising model in an external field. For a square-well symmetrical binary mixture we calculate the parameters of the critical point as functions of the microscopic parameter r measuring the relative strength of interactions between the particles of dissimilar and similar species. The calculations are performed at intermediate ($\lambda = 1.5$) and moderately long ($\lambda = 2.0$) intermolecular potential ranges. The obtained results agree well with the ones of computer simulations.

PACS numbers: 05.70.Jk

§ Author to whom any correspondence should be addressed. E-mail address: romanr@icmp.lviv.ua

1. Introduction

Binary mixtures in contrast to their constituent components can exhibit three different types of two-phase equilibria: vapour-liquid, liquid-liquid and gas-gas [1, 2]. The possibility of the realization of these phenomena and their priority depend both on the external conditions and microscopic parameters of a mixture. The study of the influence of interparticle interactions on the critical properties of a binary mixture is an interesting and actual problem. During the last decade this problem has been intensively studied by integral equation methods [3]-[9]. However, this approach, although reproduces different phase diagram types by varying the microscopic parameters, gives only a qualitative picture of the phenomena under consideration. On the other hand, the critical properties of simple fluids and binary mixtures have been studied recently using Monte Carlo (MC) simulations [10]-[14]. In [14] the vapour-liquid critical temperature was calculated for the symmetrical mixture of hard spheres interacting via the square-well potentials. Thus, it is interesting to test a theory using such a simple binary fluid model.

In the present paper we propose a microscopic approach to the study of the vapour-liquid critical point of a symmetrical binary mixture. This approach is based on the method of collective variables (CV). This method was developed in [15] and appears to be successful in describing the second order phase transition of the 3D Ising model [16] and the vapour-liquid critical point of a one-component fluid [17]. On the basis of this approach both universal and non-universal quantities were obtained.

In [18] the CV method with a reference system (RS) was generalized for the case of a grand canonical ensemble for a multicomponent continuous system. Using this approach the phase diagram of the symmetrical mixture was examined within the framework of the Gaussian approximation [19]-[21].

In this paper we determine an explicit form of the effective Ginsburg-Landau-Wilson (GLW) Hamiltonian of the symmetrical binary mixture in the vicinity of the vapour-liquid critical point. Then we integrate the functional of the grand partition function by the use of the layer-by-layer integration method proposed in [16] for the 3D Ising model. As a result of this integration one obtains recursion relations for the coefficients of the GLW Hamiltonian. The analysis of these relations yields an equation for T_c . Here we avoid extensive consideration of the results pertaining to the Ising model and call the readers' attention to [16] where this problem was studied in detail. The method which we describe here yields the same critical exponents as in [22] (see table 1).

Table 1. Values of the critical exponents and the ratios of the critical amplitudes for the 3D Ising model obtained within the framework of the CV method.

ν	α	β	γ	A^+/A^-	Γ^+/Γ^-
0.637	0.088	0.319	1.275	0.435	6.967

The paper is organized as follows. We give a functional representation of the grand partition function of a two-component continuous system in section 2 and appendix A.

In section 3 we construct the basic density measure (GLW Hamiltonian) with respect to the CV which include a variable corresponding to the order parameter. In this section we also present the basic ideas of the method of the partition function integration in the vicinity of the critical point. In section 4 we apply our formalism to calculating the critical characteristics (temperature and density) of the binary square-well symmetrical mixture. The obtained results are discussed and compared with the MC simulation data reported recently by N.B.Wilding [14] and E.de Miquel [13].

2. Functional representation of the grand partition function of a binary mixture

Let us consider a binary fluid mixture consisting of N_a particles of species "a" and N_b particles of species "b". The system is in volume V at temperature T . Let us assume that an interaction in the system has a pairwise additive character. The interaction potential between γ particle at $\mathbf{r}_i$ and δ particle at $\mathbf{r}_j$ can be expressed as a sum of two terms:

$$U_{\gamma\delta}(|\mathbf{r}_i - \mathbf{r}_j|) = \Psi_{\gamma\delta}(|\mathbf{r}_i - \mathbf{r}_j|) + \Phi_{\gamma\delta}(|\mathbf{r}_i - \mathbf{r}_j|), \quad (1)$$

where $\Psi_{\gamma\delta}(r)$ is a potential of a short-range repulsion and $\Phi_{\gamma\delta}(r)$ is an attractive part of the potential which dominates at large distances.

A functional of the grand partition function (GPF) of the binary homogeneous system in the CV method with a RS can be represented as a product of two factors (see Appendix A):

$$\Xi = \Xi_0 \Xi_1, \quad (2)$$

where Ξ_0 is the GPF of the RS which we suppose to be known. Ξ_1 is the part of the GPF which is written in the CV space:

$$\begin{aligned} \Xi_1 = \int (d\rho)(dc) \exp \left\{ \beta\mu_1^+ \rho_0 + \beta\mu_1^- c_0 - \frac{\beta}{2} \sum_{\mathbf{k}} \left[\tilde{V}(k) \rho_{\mathbf{k}} \rho_{-\mathbf{k}} + \right. \right. \\ \left. \left. + 2\tilde{U}(k) \rho_{\mathbf{k}} c_{\mathbf{k}} + \tilde{W}(k) c_{\mathbf{k}} c_{-\mathbf{k}} \right] \right\} J(\rho, c). \end{aligned} \quad (3)$$

Chemical potentials $\mu_1^+ = \frac{1}{\sqrt{2}}(\mu_1^a + \mu_1^b)$ and $\mu_1^- = \frac{1}{\sqrt{2}}(\mu_1^a - \mu_1^b)$ are determined from the conditions:

$$\frac{d \ln \Xi_1}{d \beta \mu_1^+} = \langle N_a \rangle + \langle N_b \rangle = \langle N \rangle \quad (4a)$$

$$\frac{d \ln \Xi_1}{d \beta \mu_1^-} = \langle N_a \rangle - \langle N_b \rangle. \quad (4b)$$

Functions $\tilde{V}(k)$, $\tilde{W}(k)$ and $\tilde{U}(k)$ are combinations of Fourier transforms of the initial interaction potentials $\tilde{\Phi}_{\gamma\delta}(k)$:

$$\begin{aligned}
\tilde{V}(k) &= \left(\frac{\beta^{-1}}{2}\right) [\alpha_{aa}(k) + \alpha_{bb}(k) + 2\alpha_{ab}(k)], \\
\tilde{U}(k) &= \left(\frac{\beta^{-1}}{2}\right) [\alpha_{aa}(k) - \alpha_{bb}(k)], \\
\tilde{W}(k) &= \left(\frac{\beta^{-1}}{2}\right) [\alpha_{aa}(k) + \alpha_{bb}(k) - 2\alpha_{ab}(k)].
\end{aligned} \tag{5}$$

$$\begin{aligned}
J(\rho, c) &= \int (d\nu)(d\omega) \exp \left\{ i2\pi \sum_{\mathbf{k}} (\omega_{\mathbf{k}} \rho_{\mathbf{k}} + \nu_{\mathbf{k}} c_{\mathbf{k}}) + \right. \\
&\quad \left. + \sum_{n \geq 1} \sum_{i_n \geq 0} D_n^{(i_n)}(\omega, \nu) \right\}
\end{aligned} \tag{6}$$

is a transition Jacobian to the CV $\rho_{\mathbf{k}}, c_{\mathbf{k}}$ averaged on the RS, variables $\omega_{\mathbf{k}}, \nu_{\mathbf{k}}$ are conjugated to variables $\rho_{\mathbf{k}}, c_{\mathbf{k}}$, respectively.

$$\begin{aligned}
D_n^{(i_n)}(\omega, \nu) &= \left[\frac{(-i2\pi)^n}{n!} \right] \left(\frac{1}{2} \right)^{n/2} \sum_{k_1 \dots k_n} \mathcal{M}_n^{(i_n)}(0, \dots, 0) \times \\
&\quad \times \nu_{\mathbf{k}_1} \dots \nu_{\mathbf{k}_{i_n}} \omega_{\mathbf{k}_{i_n+1}} \dots \omega_{\mathbf{k}_n} \delta_{\mathbf{k}_1 + \dots + \mathbf{k}_n}.
\end{aligned} \tag{7}$$

Index i_n indicates the number of variables $\nu_{\mathbf{k}}$ in the cumulant expansion (7). Cumulants $\mathcal{M}_n^{(i_n)}(0, \dots, 0)$ are linear combinations of the initial cumulants $\mathcal{M}_{\gamma_1 \dots \gamma_n}(0, \dots, 0)$ ($\gamma_i = a, b$) (see Appendix B).

We consider a symmetrical binary fluid mixture (SBFM), i.e. a system in which the two pure components "a" and "b" are identical and only interactions between the particles of dissimilar species differ. Notwithstanding its simplicity, the SBFM exhibits all the three types of two-phase equilibria which are observed in real binary fluids, namely: vapour-liquid, liquid-liquid and gas-gas equilibria. For the SBFM $\tilde{U}(k) = 0$ in (3) and there are only terms with even indices i_n in the cumulant expansion (7) [20].

3. The method

As it was already shown [20], the phase diagram of the SBFM consists of three ranges (see figure 1): (1) gas-gas separation and vapour-liquid phase transitions; (2) vapour-liquid and liquid-liquid phase transitions; (3) vapour-liquid phase transition only.

The order of priority of the vapour-liquid and separation phase transitions depends on both the external conditions and the microscopic properties of the system. There exist two critical temperature branches in such a system: branch (T_c^{v-l}) connected with the variable ρ_0 and branch (T_c^{sep}) connected with the variable c_0 [20]. All the thermodynamic functions of the SBFM are symmetrical with respect to the concentration $x=0.5$ and have an extremum at this point [1, 19]. The concentration $x=0.5$ is a critical one for this model mixture.




Figure 1. Three phase regions of the symmetrical mixture depending on the microscopic parameters: (1) gas-gas and vapour-liquid phase transitions ($T_c^{g-g} > T_c^{v-l}$); (2) vapour-liquid and liquid-liquid phase transitions ($T_c^{v-l} > T_c^{l-l}$); (3) vapour-liquid phase transition only. S_+ is the structure factor of the reference system.

In this paper we consider a symmetrical fluid mixture within range 2 or 3 of its phase diagram.

In the case of the SBFM the variables ρ_0 and c_0 are connected with the order parameters for the vapour-liquid and separation phase transitions, respectively [20]. This fact allows us to separate CV $\rho_{\mathbf{k}}$ and $c_{\mathbf{k}}$ into essential and non-essential ones depending on the phase transition considered. Since we are interested in the vapour-liquid critical point, we can consider CV $c_{\mathbf{k}}$ (and $\nu_{\mathbf{k}}$) to be non-essential (CV $c_{\mathbf{k}}$ do not contain a variable connected with the order parameter, the coefficients standing at the second power of $c_{\mathbf{k}}$ (and $\nu_{\mathbf{k}}$) are negative) and we can integrate over $c_{\mathbf{k}}$ (and $\nu_{\mathbf{k}}$) with the Gaussian density measure. In respect to CV $\rho_{\mathbf{k}}$ it is necessary to construct the basic density measure taking into account higher powers of $\rho_{\mathbf{k}}$ (we shall consider a ρ^4 model).

As a result of integrating in (3)-(7) over variables $c_{\mathbf{k}}$ (and $\nu_{\mathbf{k}}$) we obtain for the GPF:

$$\Xi = \Xi_0 \Xi_G^c \int (d\rho) \exp \left\{ \beta \mu_1^+ \rho_0 - \frac{\beta}{2} \sum_{\mathbf{k}} \tilde{V}(k) \rho_{\mathbf{k}} \rho_{-\mathbf{k}} \right\} J(\rho), \quad (8)$$

where

$$\Xi_G^c = \prod_{\mathbf{k}} \frac{1}{\sqrt{1 + \beta \tilde{W}(k) \mathcal{M}_2^{(2)}(0)/2}}, \quad (9)$$

$$J(\rho) = \int (d\omega) \exp \left\{ i2\pi \sum_{\mathbf{k}} \omega_{\mathbf{k}} \rho_{\mathbf{k}} + \sum_{n \geq 1} \frac{(-i2\pi)^n}{n!} \left(\frac{1}{2} \right)^{n/2} \times \right. \\ \left. \times \sum_{\mathbf{k}_1 \dots \mathbf{k}_n} \mathcal{M}_n(0) \omega_{\mathbf{k}_1} \dots \omega_{\mathbf{k}_n} \delta_{\mathbf{k}_1 + \dots + \mathbf{k}_n} \right\}, \quad (10)$$

$$\mathcal{M}_n(0) = \mathcal{M}_n^{(0)}(0) + \Delta \mathcal{M}_n. \quad (11)$$

$\Delta\mathcal{M}_n$ are the corrections obtained as the result of integration over variables $c_{\mathbf{k}}$:

$$\begin{aligned}
\Delta\mathcal{M}_1 &= \frac{\mathcal{M}_3^{(2)}(0)}{12} \frac{1}{\langle N \rangle} \sum_{\mathbf{k}} \tilde{g}(|\mathbf{k}|), \\
\Delta\mathcal{M}_2 &= \frac{\mathcal{M}_4^{(2)}(0)}{12} \frac{1}{\langle N \rangle} \sum_{\mathbf{k}} \tilde{g}(|\mathbf{k}|) + \\
&\quad + \frac{(\mathcal{M}_3^{(2)}(0))^2}{72} \frac{1}{\langle N \rangle^2} \sum_{\mathbf{k}} \tilde{g}(|\mathbf{k}|) \tilde{g}(|\mathbf{k}_1 - \mathbf{k}|), \\
\Delta\mathcal{M}_3 &= \frac{\mathcal{M}_3^{(2)}(0)\mathcal{M}_4^{(2)}(0)}{48} \frac{1}{\langle N \rangle^2} \sum_{\mathbf{k}} \tilde{g}(|\mathbf{k}|) \tilde{g}(|\mathbf{k}_1 - \mathbf{k}|) + \\
&\quad + \frac{(\mathcal{M}_3^{(2)}(0))^3}{6} \frac{1}{\langle N \rangle^3} \sum_{\mathbf{k}} \tilde{g}(|\mathbf{k}|) \tilde{g}(|\mathbf{k}_1 + \mathbf{k}|) \tilde{g}(|\mathbf{k}_2 - \mathbf{k}|), \\
\Delta\mathcal{M}_4 &= \frac{(\mathcal{M}_4^{(2)}(0))^2}{96} \frac{1}{\langle N \rangle^2} \sum_{\mathbf{k}} \tilde{g}(|\mathbf{k}|) \tilde{g}(|\mathbf{k}_1 - \mathbf{k}|) + \\
&\quad + \left(\frac{\mathcal{M}_3^{(2)}(0)}{3!}\right)^4 \frac{1}{\langle N \rangle^4} \sum_{\mathbf{k}} \tilde{g}(|\mathbf{k}|) \tilde{g}(|\mathbf{k}_1 + \mathbf{k}|) \times \\
&\quad \times \tilde{g}(|\mathbf{k}_2 - \mathbf{k}|) \tilde{g}(|\mathbf{k}_3 + \mathbf{k}_1 + \mathbf{k}|),
\end{aligned} \tag{12}$$

where

$$\tilde{g}(k) = - \frac{\beta \langle N \rangle \tilde{W}(k)}{\frac{1}{2}\beta \tilde{W}(k) \mathcal{M}_2^{(2)}(0) + 1}. \tag{13}$$

In figure 2 the typical behaviour of the potential $\tilde{V}(k)/|\tilde{V}(0)|$ is shown.




Figure 2. The behaviour of the Fourier transform $\tilde{V}(k)/|\tilde{V}(0)|$ of the attractive part of the interaction potential $V(r)$.

Let us further assume that $\tilde{V}(k) = 0$ at $|\mathbf{k}| > B$. Then, integration in (8) over $\rho_{\mathbf{k}}$ with $|\mathbf{k}| > B$ leads to δ -functions and the expression for Ξ contains only the sums over $\mathbf{k}$ with $|\mathbf{k}| \leq B$.

We consider a set of $\mathbf{k}$ vectors, $|\mathbf{k}| \leq B$, as corresponding to the sites of a reciprocal lattice conjugated to a certain block lattice $\{r_l\}$ with N_B block sites in the periodicity volume V :

$$\langle N_B \rangle = \frac{V}{C^3} = \frac{V}{(\pi/B)^3} = \frac{(B\sigma)^3 \langle N \rangle}{6\pi^2\eta}, \quad (14)$$

$\eta = \frac{\pi}{6}\rho\sigma^3$ is fraction density. Therefore, one may consider quantity B as the size of the first Brillouin zone of this block lattice.

The shift

$$\begin{aligned} \omega_{\mathbf{k}} &= \omega'_{\mathbf{k}} + \Delta\delta_{\mathbf{k}}, \\ \rho_{\mathbf{k}} &= \rho'_{\mathbf{k}} + \tilde{\mathcal{M}}_1\delta_{\mathbf{k}}, \end{aligned}$$

where

$$\Delta = -\frac{i}{2\pi} \frac{\bar{\mathcal{M}}_3(0)}{\bar{\mathcal{M}}_4(0)} \delta_{\mathbf{k}},$$

$$\tilde{\mathcal{M}}_1 = \bar{\mathcal{M}}_1(0) - \frac{\bar{\mathcal{M}}_2(0)\bar{\mathcal{M}}_3(0)}{\bar{\mathcal{M}}_4(0)} + \frac{\bar{\mathcal{M}}_3^3(0)}{3\bar{\mathcal{M}}_4^2(0)},$$

$(\bar{\mathcal{M}}_n(0) = \frac{\mathcal{M}_n(0)}{(\sqrt{2})^n}, n = 1, \dots, 4)$, transforms Ξ into a form containing terms $\tilde{\mathcal{M}}_1(0)$, $\tilde{\mathcal{M}}_2(0)$ and $\tilde{\mathcal{M}}_4(0)$ only (the primes on $\rho_{\mathbf{k}}$ and $\omega_{\mathbf{k}}$ are omitted for clarity):

$$\begin{aligned} \Xi &= \Xi_0 \Xi_G^{(1)} \int \exp \left\{ \mu^* \rho_0 - \frac{\beta}{2} \sum_{\mathbf{k} < B} \tilde{V}(k) \rho_{\mathbf{k}} \rho_{-\mathbf{k}} + \right. \\ &\quad \left. + i2\pi \sum_{\mathbf{k} < B} \omega_{\mathbf{k}} \rho_{\mathbf{k}} - \frac{(2\pi)^2}{2} \tilde{\mathcal{M}}_2(0) \sum_{\mathbf{k} < B} \omega_{\mathbf{k}} \omega_{-\mathbf{k}} - \frac{(2\pi)^4}{4! \langle N_B \rangle} |\tilde{\mathcal{M}}_4(0)| \times \right. \\ &\quad \left. \times \sum_{\mathbf{k}_1 \dots \mathbf{k}_4 < B} \omega_{\mathbf{k}_1} \omega_{\mathbf{k}_2} \omega_{\mathbf{k}_3} \omega_{\mathbf{k}_4} \delta_{\mathbf{k}_1 + \dots + \mathbf{k}_4} \right\} (d\omega)^{N_B} (d\rho)^{N_B}, \end{aligned} \quad (15)$$

where

$$\begin{aligned} \Xi_G^{(1)} &= \Xi_G^c \exp \left\{ \mu^* \tilde{\mathcal{M}}_1 + \frac{\beta \tilde{V}^*(0)}{2} \tilde{\mathcal{M}}_1^2 - \right. \\ &\quad \left. - \frac{\bar{\mathcal{M}}_1(0)\bar{\mathcal{M}}_3(0)}{\bar{\mathcal{M}}_4(0)} - \frac{\bar{\mathcal{M}}_2(0)\bar{\mathcal{M}}_3^2(0)}{2\bar{\mathcal{M}}_4^2(0)} - \frac{\bar{\mathcal{M}}_3^4(0)}{8\bar{\mathcal{M}}_4^3(0)} \right\}, \\ \mu^* &= h - a_1, \quad a_1 = \frac{\bar{\mathcal{M}}_3(0)}{|\bar{\mathcal{M}}_4(0)|} + \beta \tilde{V}^*(0) \tilde{\mathcal{M}}_1, \quad h = \beta \mu_1^+, \\ \tilde{\mathcal{M}}_2(0) &= \bar{\mathcal{M}}_2(0) - \frac{\bar{\mathcal{M}}_3^2(0)}{2\bar{\mathcal{M}}_4(0)}, \\ \tilde{\mathcal{M}}_4(0) &= \langle N_B \rangle \bar{\mathcal{M}}_4(0). \end{aligned} \quad (16)$$

Expression (15) for Ξ corresponds to the Ising model in the external field $(a_1 - \beta \mu_1^+)$ with the only difference: cumulants $\tilde{\mathcal{M}}_2(0)$, $\tilde{\mathcal{M}}_4(0)$ are functions of the fraction density η , temperature T and parameters of the attractive interaction $\tilde{\Phi}_{\gamma\delta}(k)$.

After integration over $\omega_{\mathbf{k}}$ we obtain the following form for the GPF:

$$\begin{aligned} \Xi &= \Xi_0 \Xi_G^{(1)} \left[Z(\tilde{\mathcal{M}}_2, \tilde{\mathcal{M}}_4) \right]^{<N_B>} (\sqrt{2})^{<N_B>-1} \times \\ &\times \int \exp[E_4(\rho)] (d\rho)^{N_B}. \end{aligned} \quad (17)$$

Here

$$\begin{aligned} E_4(\rho) &= \mu^* \rho_0 - \frac{1}{2} \sum_{\mathbf{k} < B} d_2(k) \rho_{\mathbf{k}} \rho_{-\mathbf{k}} - \frac{a_4}{4! <N_B>} \times \\ &\times \sum_{\mathbf{k}_1 \dots \mathbf{k}_4 < B} \rho_{\mathbf{k}_1} \dots \rho_{\mathbf{k}_4} \delta_{\mathbf{k}_1 + \dots + \mathbf{k}_4} + \dots, \end{aligned} \quad (18)$$

$$\begin{aligned} Z(\tilde{\mathcal{M}}_2, \tilde{\mathcal{M}}_4) &= \left(\frac{1}{2\pi} \right)^{1/2} \left(\frac{3}{|\tilde{\mathcal{M}}_4(0)|} \right)^{1/4} e^{x^2/4} U(0, x), \\ d_2(k) &= a_2 + \beta \tilde{V}(k), \quad a_2 = \sqrt{\frac{3}{|\tilde{\mathcal{M}}_4(0)|}} K(x), \\ a_4 &= \frac{3}{|\tilde{\mathcal{M}}_4(0)|} L(x), \end{aligned} \quad (19)$$

where

$$\begin{aligned} K(x) &= U(1, x)/U(0, x), \\ L(x) &= 3K^2(x) + 2xK(x) - 2, \\ x &= \sqrt{\frac{3}{|\tilde{\mathcal{M}}_4(0)|}} \tilde{\mathcal{M}}_2(0). \end{aligned} \quad (20)$$

$U(a, x)$ is a parabolic cylinder function [23]. Expressions (17)-(20) have the same forms as similar expressions for a one-component system obtained in [17]. This coincidence is achieved due to the symmetry of the model under consideration. $E_4(\rho)$ is the Ginzburg-Landau-Wilson Hamiltonian for the SBFM in the vicinity of the vapour-liquid critical point.

In order to integrate the GPF (17)-(20) over $\rho_{\mathbf{k}}$ and determine the critical temperature we use the method developed in [16, 24] for the Ising model. The essence of the method consists in subsequent integration over the layers of the CV space, beginning from $\rho_{\mathbf{k}}$ which correspond to short-wave fluctuations. Variations of the coefficients of $E_4(\rho)$ as the result of integration over $\rho_{\mathbf{k}}$ in n subsequent layers of CV phase space are described by the recursion formulae derived in [16]. For the case $T > T_c$ in the interval $[0, B]$ there exist three characteristic regions [16]. The first region $B_{m_\tau} < k \leq B$ corresponds to the strongly correlated fluctuations $\rho_{\mathbf{k}}$, their density measure is non-Gaussian. The procedure based on the renormalization group symmetry is valid here. This is the region of the critical regime (CR). The second region $0 < k \leq B_{m_\tau}$ is related to the fluctuation distributed according to the Gaussian density measure. This is the limiting Gaussian regime (LGR).

The third region consists of the point $k = 0$. The variable ρ_0 is a macroscopic one and corresponds to the fluctuations of the particle density in the "external field" μ^* .

We integrate (17) according to the following scheme [16]. The region $(0, B)$ is divided into the intervals $(B_1, B), \dots, (B_{i+1}, B_i), \dots$, where $B_n = B/S^n$ (S is a division parameter). Each interval corresponds to a layer of subscripts $\mathbf{k}$ in the Brillouin zone and each layer of subscripts $\mathbf{k}$ - to a layer in the phase space $\rho_{\mathbf{k}}$. Integrating gradually over the layers we get a block lattice sequence with an appropriately growing block period and with the Hamiltonian corresponding to each block. Each Hamiltonian is characterized by the coefficients $d_2, a_4; d_2^{(1)}, a_4^{(1)}; d_2^{(2)}, a_4^{(2)}$, etc.. For the sequence of the block Hamiltonians $\{d_2^{(n)}, a_4^{(n)}\}$ the renormalization group symmetry holds and the fixed point is of a saddle type. Because the explicit expressions for the initial values of coefficients $d(k)$ and a_4 are given (see (18)-(20)), the solutions of the renormalization group type are functions of microscopic parameters, density and temperature.

The CR takes place for all the variables $\rho_{\mathbf{k}}$ at the critical point. Therefore, the critical temperature can be determined from the solution of recurrent equations. Using the result obtained in [24] we derive the formula

$$A(\beta\tilde{V}(0))^2 + B(\beta\tilde{V}(0)) + D = 0, \quad (21)$$

where

$$\begin{aligned} A &= 1 - f_0 - R^{(0)}\sqrt{\varphi_0}, \\ B &= -a_2, \\ D &= a_4 R^{(0)} / \sqrt{\varphi_0}. \end{aligned}$$

f_0, φ_0 are coordinates of the reduced fixed point, $R^{(0)}$ is a universal function of parameter S . The optimal value of S is 3.4252 and the values of $f_0, \varphi_0, R^{(0)}$ corresponding to it are taken from [24]. From the condition $\mu^* = 0$ we obtain the second equation [17]:

$$\mathcal{M}_3(0) = 0, \quad (22)$$

which allows us to determine the critical density of the system.

4. Results and discussions

In this section we present our results for the vapour-liquid critical point of symmetrical mixtures, using the method proposed above. These results are compared with those previously obtained by Monte-Carlo simulations [13, 14].

The system under study is a symmetrical hard sphere square-well binary mixture. The interaction potential between the particles is given by:

$$U_{\gamma\delta}(r) = \begin{cases} \infty, & \text{if } r < \sigma \\ -\epsilon_{\gamma\delta}, & \text{if } \sigma \leq r < \lambda\sigma \\ 0, & \text{if } r \geq \lambda\sigma \end{cases},$$

where σ is a hard sphere diameter, λ is a range of the potential, and $\epsilon_{\gamma\delta}$ is a well-depth of the interaction between the particles of types γ and δ . For a symmetrical mixture

$\epsilon_{aa} = \epsilon_{bb} = \epsilon \neq \epsilon_{ab}$. In our formalism a completely analytical treatment for general λ is possible. But in this paper we choose $\lambda = 1.5$ and $\lambda = 2.0$, because for these values of λ the MC results are available.

We split the potential $U_{\gamma\delta}(r)$ into short- and long-range parts using the Weeks-Chandler-Andersen partition [27]. As a result, we have:

$$\Psi_{\gamma\delta}(r) = \begin{cases} \infty, & r \leq \sigma \\ 0, & r > \sigma \end{cases}, \quad (23)$$

$$\Phi_{\gamma\delta}(r) = \begin{cases} -\epsilon_{\gamma\delta}, & 0 \leq r \leq \lambda\sigma \\ 0, & r > \lambda\sigma \end{cases}. \quad (24)$$

In our case the RS is a hard sphere system with the diameter σ (see (23)). The Fourier transform of function (24) has the form:

$$\tilde{\Phi}_{\gamma\delta}(k) = \tilde{\Phi}_{\gamma\delta}(0) \frac{3}{(\lambda x)^3} [-\lambda x \cos(\lambda x) + \sin(\lambda x)],$$

where

$$x = k\sigma, \\ \tilde{\Phi}_{\gamma\delta}(0) = -\epsilon_{\gamma\delta}\sigma^3 \frac{4\pi}{3} \lambda^3.$$

Cumulants $\mathcal{M}_n^{(in)}(0, \dots, 0)$ are calculated according to the formulae given in Appendix B. For $S_2(0)$ the Percus-Yevick approximation is used:

$$S_2(0) = \frac{(1 - \eta)^4}{(1 + 2\eta)^2}.$$

The solutions of equations (21)-(22) are found numerically using a self-consistent procedure by means of which the dependences of the coefficients a_2 and a_4 (as well as cumulants $\mathcal{M}_n(0)$) on β_c are taken into consideration.

The vapour-liquid critical temperatures T_c ($T_c = k_B T / \epsilon$) versus the microscopic parameter r ($r = \epsilon_{ab} / \epsilon$ is a dissimilar interaction strength) are shown for $\lambda = 1.5$ and $\lambda = 2.0$ in figure 3.

In figure 4 we demonstrate the dependence of the critical density η_c on r for $\lambda=1.5$ and $\lambda=2.0$. We compare our results with those obtained from MC simulations: for $\lambda = 1.5$ and $r=0.72$ we have $T_c = 1.055$, while the MC simulations give $T_c = 1.06(1)$ [14], for $\lambda = 2.0$ and $r=1.0$ (the case of $r=1.0$ corresponds to a one-component system) we obtain $T_c = 2.753$ and $\eta_c = 0.129$, while the simulations give $T_c = 2.684(51)$ and $\eta_c = 0.123(43)$ [13].

5. Conclusions

In this paper we propose a method for the study of the vapour-liquid critical point of a symmetrical binary mixture depending on its microscopic properties. Using this method we calculated the critical temperature and critical density of the symmetrical




Figure 3. The vapour-liquid critical temperature as a function of the microscopic parameter r at $\lambda = 1.5$ (left) and $\lambda = 2.0$ (right).




Figure 4. The critical density as a function of the microscopic parameter r .

hard sphere square-well fluid versus the microscopic parameter r which is the measure of the dissimilar interaction in the system. Our results agree with those obtained by MC simulations. We can improve our results in the following ways: 1) taking into consideration the region of $\mathbf{k}$ with $|\mathbf{k}| > B$ (see figure 2); 2) using a higher approximation than the ρ^4 one.

Appendix A

A grand partition function of a two-component fluid system in the CV representation with a RS can be written as in [18]:

$$\Xi = \Xi_0 \Xi_1,$$

where

$$\Xi_0 = \sum_{N_a=0}^{\infty} \sum_{N_b=0}^{\infty} \prod_{\gamma=a}^b \exp \left[\frac{\beta \mu_0^\gamma N_\gamma}{N_\gamma!} \right] \int (d\Gamma) \exp \left[-\frac{\beta}{2} \sum_{\gamma,\delta=a,b} \sum_{i,j} \psi_{\gamma\delta}(r_{ij}) \right]$$

is a grand partition function of the RS; $\beta = \frac{1}{k_B T}$, k_B is the Boltzman constant, T is temperature; $(d\Gamma) = \prod_{a,b} d\Gamma_{N_\gamma}$, $d\Gamma_{N_\gamma} = d\vec{r}_1^\gamma d\vec{r}_2^\gamma \dots d\vec{r}_{N_\gamma}^\gamma$ is a volume element of the configurational space of the γ -th species; μ_0^γ is a chemical potential of the γ -th species in the RS.

The part of the grand partition function which is defined in the CV phase space has the form of a functional integral:

$$\Xi_1 = \int (d\rho) \exp \left[\beta \sum_{\gamma} \mu_1^\gamma \rho_{0,\gamma} - \frac{1}{2\beta} \sum_{\gamma,\delta=a,b} \sum_{\mathbf{k}} \alpha_{\gamma\delta}(k) \rho_{\mathbf{k},\gamma} \rho_{-\mathbf{k},\delta} \right] J(\rho_a, \rho_b). \quad (\text{A.1})$$

Here,

1) μ_1^γ is a part of the chemical potential of the γ -th species

$$\mu_1^\gamma = \mu_\gamma - \mu_0^\gamma + \frac{1}{2\beta} \sum_{\mathbf{k}} \alpha_{\gamma\gamma}(k),$$

and is determined from the equation

$$\frac{\partial \ln \Xi_1}{\partial \beta \mu_1^\gamma} = \langle N_\gamma \rangle,$$

μ_γ is a full chemical potential of the γ -th species; $\alpha_{\gamma\delta}(k) = \frac{\beta}{V} \tilde{\phi}_{\gamma\delta}(k)$; $\langle N_\gamma \rangle$ is an average number of the γ -th species particles.

2) $\rho_{\mathbf{k},\gamma} = \rho_{\mathbf{k},\gamma}^c - i\rho_{\mathbf{k},\gamma}^s$ ($\gamma = a, b$) are collective variables of the γ -th species, the indices c and s denote the real part and the coefficient at the imaginary part of $\rho_{\mathbf{k},\gamma}$; $\rho_{\mathbf{k},\gamma}^c$ and $\rho_{\mathbf{k},\gamma}^s$ describe the value of the $\mathbf{k}$ -th fluctuation mode of the number of the γ -th species particles. Each $\rho_{\mathbf{k},\gamma}^c$ and $\rho_{\mathbf{k},\gamma}^s$ takes all the real values from $-\infty$ to $+\infty$. $(d\rho)$ is a volume element of the CV phase space:

$$(d\rho) = \prod_{\gamma} d\rho_{0,\gamma} \prod'_{\mathbf{k} \neq 0} d\rho_{\mathbf{k},\gamma}^c d\rho_{\mathbf{k},\gamma}^s.$$

The prime means that the product over $\mathbf{k}$ is performed in the upper semi-space;

3) $J(\rho_a, \rho_b)$ is a transition Jacobian to the CV averaged on the RS:

$$J(\rho_a, \rho_b) = \int (d\nu) \prod_{\gamma=a}^b \exp \left[i2\pi \sum_{\vec{k}} \nu_{\mathbf{k},\gamma} \rho_{\mathbf{k},\gamma} \right] \exp \left[\sum_{n \geq 1} \frac{(-i2\pi)^n}{n!} \times \sum_{\gamma_1 \dots \gamma_n} \sum_{\mathbf{k}_1 \dots \mathbf{k}_n} M_{\gamma_1 \dots \gamma_n}(\mathbf{k}_1, \dots, \mathbf{k}_n) \nu_{\mathbf{k}_1, \gamma_1} \dots \nu_{\mathbf{k}_n, \gamma_n} \right],$$

where variables $\nu_{\mathbf{k},\gamma}$ are conjugated to CV $\rho_{\mathbf{k},\gamma}$. $M_{\gamma_1 \dots \gamma_n}(\mathbf{k}_1, \dots, \mathbf{k}_n)$ is the n -th cumulant connected with $S_{\gamma_1 \dots \gamma_n}(k_1, \dots, k_n)$, the n -particle partial structure factor of the RS, by means of the relation

$$M_{\gamma_1 \dots \gamma_n}(\mathbf{k}_1, \dots, \vec{k}_n) = \sqrt[n]{N_{\gamma_1} \dots N_{\gamma_n}} S_{\gamma_1 \dots \gamma_n}(k_1, \dots, k_n) \delta_{\mathbf{k}_1 + \dots + \mathbf{k}_n},$$

where $\delta_{\mathbf{k}_1+\dots+\mathbf{k}_n}$ is a Kronecker symbol.

In general, the dependence of $M_{\gamma_1\dots\gamma_n}(\mathbf{k}_1, \dots, \mathbf{k}_n)$ on wave vectors $\mathbf{k}_1, \dots, \mathbf{k}_n$ is complicated. Hereafter we shall replace $M_{\gamma_1\dots\gamma_n}(\mathbf{k}_1, \dots, \mathbf{k}_n)$ by their values in the long-wave length limit $M_{\gamma_1\dots\gamma_n}(0, \dots, 0)$;

4) $\tilde{\phi}_{\gamma\delta}(k)$ is a Fourier transform of the attractive potential $\phi_{\gamma\delta}(r)$. Function $\tilde{\phi}_{\gamma\delta}(k)$ satisfies the following requirements: $\tilde{\phi}_{\gamma\delta}(k)$ is negative for small values of $\mathbf{k}$ and $\lim_{\mathbf{k} \rightarrow \infty} \tilde{\phi}_{\gamma\delta}(k) = 0$.

We pass in (A.1) to CV $\rho_{\mathbf{k}}$ and $c_{\mathbf{k}}$ (according to $\omega_{\mathbf{k}}$ and $\gamma_{\mathbf{k}}$) by means of the orthogonal linear transformation:

$$\begin{aligned}\rho_{\mathbf{k}} &= \frac{\sqrt{2}}{2}(\rho_{\mathbf{k},a} + \rho_{\mathbf{k},b}), & c_{\mathbf{k}} &= \frac{\sqrt{2}}{2}(\rho_{\mathbf{k},a} - \rho_{\mathbf{k},b}), \\ \omega_{\mathbf{k}} &= \frac{\sqrt{2}}{2}(\nu_{\mathbf{k},a} + \nu_{\mathbf{k},b}), & \nu_{\mathbf{k}} &= \frac{\sqrt{2}}{2}(\nu_{\mathbf{k},a} - \nu_{\mathbf{k},b}).\end{aligned}\tag{A.2}$$

Now $\rho_{\mathbf{k}}$ and $c_{\mathbf{k}}$ are connected with the total density fluctuation modes and the relative density (or concentration) fluctuation modes, respectively.

As a result, for Ξ_1 we obtain formulae (3)-(7).

Appendix B

Cumulants $\mathcal{M}_n^{(i_n)}(0)$ with $n \leq 4$ are expressed in terms of the initial cumulants $\mathcal{M}_{\gamma_1\dots\gamma_n}(0, \dots, 0)$ ($\gamma_1, \dots, \gamma_n = a, b$) as follows [20]:

$$\begin{aligned}\mathcal{M}_1^{(0)}(0) &= \mathcal{M}_a(0) + \mathcal{M}_b(0) = \langle N \rangle \\ \mathcal{M}_1^{(1)}(0) &= \mathcal{M}_a(0) - \mathcal{M}_b(0) = \langle N_a \rangle - \langle N_b \rangle \\ \mathcal{M}_2^{(0)}(0) &= \mathcal{M}_{aa}(0) + \mathcal{M}_{bb}(0) + 2\mathcal{M}_{ab}(0) \\ \mathcal{M}_2^{(1)}(0) &= \mathcal{M}_{aa}(0) - \mathcal{M}_{bb}(0) \\ \mathcal{M}_2^{(2)}(0) &= \mathcal{M}_{aa}(0) + \mathcal{M}_{bb}(0) - 2\mathcal{M}_{ab}(0) \\ \mathcal{M}_3^{(0)}(0) &= \mathcal{M}_{aaa}(0) + \mathcal{M}_{bbb}(0) + 3[\mathcal{M}_{aab}(0) + \mathcal{M}_{abb}(0)] \\ \mathcal{M}_3^{(1)}(0) &= \mathcal{M}_{aaa}(0) - \mathcal{M}_{bbb}(0) + \mathcal{M}_{aab}(0) - \mathcal{M}_{abb}(0) \\ \mathcal{M}_3^{(2)}(0) &= \mathcal{M}_{aaa}(0) + \mathcal{M}_{bbb}(0) - \mathcal{M}_{aab}(0) - \mathcal{M}_{abb}(0) \\ \mathcal{M}_3^{(3)}(0) &= \mathcal{M}_{aaa}(0) - \mathcal{M}_{bbb}(0) - 3[\mathcal{M}_{aab}(0) - \mathcal{M}_{abb}(0)] \\ \mathcal{M}_4^{(0)}(0) &= \mathcal{M}_{aaaa}(0) + \mathcal{M}_{bbbb}(0) + \\ &\quad + 4[\mathcal{M}_{aaab}(0) + \mathcal{M}_{abbb}(0)] + 6\mathcal{M}_{aabb}(0) \\ \mathcal{M}_4^{(1)}(0) &= \mathcal{M}_{aaaa}(0) - \mathcal{M}_{bbbb}(0) + 2[\mathcal{M}_{aaab}(0) - \mathcal{M}_{abbb}(0)] \\ \mathcal{M}_4^{(2)}(0) &= \mathcal{M}_{aaaa}(0) + \mathcal{M}_{bbbb}(0) - 2\mathcal{M}_{aabb}(0) \\ \mathcal{M}_4^{(3)}(0) &= \mathcal{M}_{aaaa}(0) - \mathcal{M}_{bbbb}(0) - 2[\mathcal{M}_{aaab}(0) - \mathcal{M}_{abbb}(0)] \\ \mathcal{M}_4^{(4)}(0) &= \mathcal{M}_{aaaa}(0) + \mathcal{M}_{bbbb}(0) - \\ &\quad - 4[\mathcal{M}_{aaab}(0) + \mathcal{M}_{abbb}(0)] + 6\mathcal{M}_{aabb}(0)\end{aligned}\tag{B.1}$$

The same expressions hold at $\mathbf{k}_i \neq 0$.

The n th cumulant $\mathcal{M}_n^{(i_n)}(0)$ with $i_n = 0$ is connected with the n th structure factor of the one-component system $S_n(0)$ [20]:

$$\mathcal{M}_n^{(0)}(0) = \langle N \rangle S_n(0).$$

Structure factors $S_n(0)$ ($n \geq 2$) can be obtained from $S_2(0)$ by means of a chain of equations for correlation functions [26]. Cumulants with $i_n \neq 0$ can be expressed in terms of $\mathcal{M}_n^{(0)}(0)$ (see formulae (4.8) in [20]).

References

- [1] Van Konynenburg P H and Scott R L 1980 *Phyl. Trans. Ray. Soc.* **298A** 1442
- [2] Rowlinson J S and Swinton F L 1982 *Liquids and Liquid Mixtures* (London: Butterworth)
- [3] Abramo M C, Caccamo C and Giunta G 1986 *Phys. Rev.* **A34** 3279
- [4] McGuigan D B and Monson P A 1987 *Mol. Phys.* **62** 3
- [5] Caccamo C and Malescio G 1989 *Phys. Rev.* **A40** 6384
- [6] Malescio G 1990 *Phys. Rev.* **A42** 6241
- [7] Chen X S and Forstmann F 1992 *J. Chem. Phys.* **97** 3696
- [8] Malescio G 1992 *J. Chem. Phys.* **96** 648
- [9] Caccamo C and Giunta G 1993 *Mol. Phys.* **78** 83
- [10] Recht J R and Panagiotopoulos A Z 1993 *Mol. Phys.* **80** 843
- [11] Green D G, Jackson G, de Miguel E and Rull L R 1994 *J. Chem. Phys.* **101** 3190
- [12] Wilding N B 1997 *Phys. Rev.* **E55** 6627
- [13] de Miguel E 1997 *Phys. Rev.* **E55** 1347
- [14] Wilding N B 1998 *Phys. Rev.* **E58** 2201
- [15] Yukhnovskii I R and Holovko M F 1980 *Statistical Theory of Classic Equilibrium Systems* (Kyiv: Naukova dumka) (in Russian).
- [16] Yukhnovskii I R 1987 *Phase Transitions of the Second Order. The Collective Variables Method*, (Singapore: World Scientific)
- [17] Yukhnovskii I R, Kolomiets V O and Idzyk I M 1995 *Journal of Stat. Phys.* **80** 405
- [18] Patsahan O V and Yukhnovskii I R 1990 *Theor. Math. Phys.* **83** 387
- [19] Patsahan O V 1995 *Cond. Matt. Phys.* No.5 124
- [20] Yukhnovskii I R and Patsahan O V 1995 *Journal of Stat. Phys.* **81** 647
- [21] Patsahan O V 1996 *Ukr. Fiz. Zhurn.* **41** 877 (in Ukrainian)
- [22] Kozlovskii M P, Pylyuk I V and Duhovii V V 1997 *Cond. Mat. Phys.* No.11 17
- [23] Abramovitz M and Stegun I A 1979 *Handbook of mathematical functions* (Moscow: Nauka) (in Russian)
- [24] Kozlovskii M P and Yukhnovskii I R 1997 *ICMP-97-25U preprint* (in Ukrainian)
- [25] Yukhnovskii I R and Idzyk I M 1985 *ITF-85-97R preprint* (in Russian)
- [26] Stell G 1964 *In the equilibrium theory of classical fluids* (New York: Benjamin)
- [27] Weeks J D, Chandler D and Andersen H C 1971 *J. Chem. Phys.* **54** 12

Figure captions

Figure 1. Three phase regions of the symmetrical mixture depending on the microscopic parameters: (1) gas-gas and vapour-liquid phase transitions ($T_c^{g-g} > T_c^{v-l}$); (2) vapour-liquid and liquid-liquid phase transitions ($T_c^{v-l} > T_c^{l-l}$); (3) vapour-liquid phase transition only. S_+ is the structure factor of the reference system.

Figure 2. The behaviour of the Fourier transform $\tilde{V}(k)/|\tilde{V}(0)|$ of the attractive part of the interaction potential $V(r)$.

Figure 3. The vapour-liquid critical temperature as a function of the microscopic parameter r at $\lambda = 1.5$ (left) and $\lambda = 2.0$ (right).

Figure 4. The critical density as a function of the microscopic parameter r .